

UNIVERSITY OF GONDAR
COLLEGE OF NATURAL AND COMPUTATIONAL SCIENCES
DEPARTMENT OF APPLIED MICRO BIOLOGY



**EVALUATION OF ANTAGONIST FUNGI FOR THE BIOLOGICAL
CONTROL OF *EICHHORNIA CRASSIPES* FROM LAKE TANA**

A thesis submitted to the Department of Biology, College of Natural and Computational Sciences, University of Gondar in partial fulfillment of the requirements for the degree of Master of Sciences in Applied Microbiology.

BY: Erehemet Belete

ADVISOR: Dr. Samuel Sahile (BSc, MSc., PhD)

June, 2016
Gondar, Ethiopia

UNIVERSITY OF GONDAR
COLLEGE OF NATURAL AND COMPUTATIONAL SCIENCES
DEPARTMENT OF APPLIED MICRO BIOLOGY

A thesis submitted to the Department of Biology, Natural and Computational Sciences, University of Gondar in partial fulfillment of the requirements for the degree of Master of Sciences in Applied Microbiology.

Name of Student	Erehemet Belete (BSc.)
Name of Advisor	Dr. Samuel Sahile (BSc, MSc., PhD)
Full title of the research project	Evaluation of antagonist fungi for the biological control of <i>Eichhornia crassipes</i> in lake Tana.

Advisor's Name	Signature	Date
_____	_____	_____

Examiner's Name	Signature	Date
_____	_____	_____
_____	_____	_____

Declaration

I, the undersigned, declare that this scientific research thesis is my own original work. We agree to accept responsibility for the scientific ethical and technical conduct of the research and for the provision of required progress reports as per terms and conditions of the Research. All sources of materials and ideas for this proposal has been duly acknowledged.

Name of the Student with signature and Date

Name Erehemet Belete Signature_____ Date_____

Name of Advisor with signature and Date

Name Dr. Samuel Sahile Signature_____ Date_____

ACKNOWLEDGEMENTS

First of all I would like to thank Allah for giving me the strength and motivation through the entire period of this study to undertake and accomplish this work.

I am grateful to my advisor, Dr. Samuel Sahile, for his constructive advices and critical comments from the beginning to the end of the research work. His valuable advice and logical comments helped me a great deal to shape the paper in its present form. His fatherly approach and financial support also played a key role for the successful accomplishment of the paper. Without his help, this paper could have not taken its present form.

My heartfelt gratitude shall go to my husband Yesuf Nurhusien for his endurance and priceless suggestion he gave me throughout the course of the study. In addition, deeply felt appreciation goes to Ebrahim Nurhusein and my friends who was in one way or another contributed to the accomplishment of this work.

I would like to give special thanks to my brothers and sisters for all the assistance they have made during my early school days and subsequently.

Finally, I would like to extend my sincere thanks to microbiology laboratory assistance Adugna Assefa, Instructor Endalkachew Bizualem and the partially financial support for labour, chemical and Petri plate purchasing from EEFRI(Ethiopian Environmental and Forest research Institute).

I thank Department of Biology Gondar University for the opportunity and conduct this study and for allowing me to use the laboratory facilities and chemicals throughout the course of the study

Table of content

Contents	Pages
List of abbreviations.....	i
List of table.....	ii
Abstract.....	iii
1.Introduction.....	1
2. Review of literature.....	5
2.1 The plant water hyacinth <i>E. craccsipes</i>	5
2.1.1 Habitat.....	5
2.2 Origin of water hyacinth.....	7
2.3 Importance of water hyacinth.....	7
2.4 Water hyacinth and threats	8
2.4.1 Threats posed by water hyacinth.....	9
2.4.1.1 Destruction of biodiversity.....	9
2.4.1.2 Oxygen depletion and reduced water quality fisheries.....	9
2.4.1.3 Breeding ground for pests and vector.....	10
2.4.1.4 Blockage of waterways.....	10
2.5 Economical impact of water hyacinth.....	11
2.6 Characteristics of water hyacinth.....	12
2.6.1 Chemical composition.....	12
2.7 Management option of water hyacinth.....	13
2.7.1 Chemical control	14
2.7.2 Mechanical control	15
2.7.3 Biological control	16
2.7.3.1 Biological control of aquatic weeds with plant pathogen.....	16
2.7.4 Reduction by utilization	19
Statement of the problem	19
Significance of the study	20

General objective	21
Specific objective.....	21
3. Materials and method	21
3.1 Study area.....	21
3.2 Study design.....	21
3.3 Sampling method	21
3.4 Sampling procedures and isolation methods.....	22
3.5 Isolation and purification of pathogens from soil.....	22
3.5.1 Pathogen city test for fungal isolates on water hyacinth.....	23
3.5.2 Whole plant pathogen city test.....	23
3.6 Data source and data collection methods.....	24
3.7 Data analysis	24
3.8 Quality control	24
4.Result.....	25
4.1 Pathogenic effects of selected fungal species on <i>E. crassips</i>	25
4.1.1 Lesion size.....	25
4.1.2 Characteristics of selected fungi.....	26
4.2 Disease incidence.....	27
4.3 Disease severity.....	28
5. Discussion.....	30
6. Conclusion.....	32
7. Recommendation.....	33
8. References.....	34
Appendix.....	45

LIST OF ABBREVIATIONS

IUCNS	International Union for Conservation of nature
PDA	Potato Dextrose Agar
GEF	Global Environment Facility
DI	Disease Incidence
DS	Disease Severity
EEFRI	Ethiopian Environmental and Forest research institute
DAS	Development Actors
MEA	Malt Extract Agar
NB	Nutrient Broth
MEB	Malt Extract Broth

List of table**Pages**

Table 1 Mean lesion size caused by different fungi on test water hyacinth leaves under laboratory conditions.....	26
Table 2 Label of disease intensity effect of fungal isolates on water hyacinth under pot experiment condition.....	27
Table 3 Mean disease incidence caused by selected fungal.....	28
Table 4 Disease severity of pathogen according to Freeman and Chardatta disease severity rating scale method.....	29

ABSTRACT

Water Hyacinth (*Eichhornia crassipes* Martius) is a freshwater aquatic plant, belonging to the family Pontederiaceae and is a native of Brazil and Ecuador region. Water hyacinth often invades bodies of water that have been impacted by human activities. The aims of this study was to isolate, screen and rank fungi infecting water hyacinth plants on the basis of their pathogenicity and ability to cause damage, and to test the potential of highly pathogenic species as biocontrol agents. The experimental design was laboratory based experiment. Samples for characterization and isolation of pathogenic species of fungi, infected parts (leaf and petiole) were collected from the area, in clean plastic bags, and stored in a refrigerator at 4⁰ C. Samples were randomly selected from Lake Tana specifically Lemba, Gorgora and from the soil in microbiology laboratory. Different fungal antagonistic species were isolated from surface sterilized *Eichhornia crassipes* parts with pronounced blight syndrome. The most common species were characterized, identified and tested. A total of 12 fungi isolates potentially suspected to be biocontrols these are, *Tricothecium roseum*, *Trichoderma* spp1 and 2, *Rhizoctonia* spp, *coletricum germinal*, *Fusarium* spp, *Scieroticum* spp, *Rhizoctonia botaticola*, *Alternaria solani*, *Aspergillus flaves*, *Aspergillus niger* and *Botrytus fabae*. From detached leaf test based on the width of the lesion, the isolated species were divided into highly pathogenic, pathogenic and non-pathogenic. Of these species only highly pathogenic species were used for whole plant photogenic test. The disease incidence (DI) was taken as the percentage number of leaves on the plant that exhibited disease symptoms. The highest DI was recorded in *Rhizoctonia* spp(66%)secondly *Ticothecium roseum*(50.7%) followed by *Aspergillus flaves*(49.1%) *Fusarium* spp(43.3%) and *Aspergillus niger*(42.6%). Disease severity (DS) was assessed using the numerical rating scale. Of these species only *Rhizoctonia* spp (100%) and *Fusarium* spp (78.6%) showed high disease severity. These fungi were associated with a high percentage of tissue death after five weeks application. *Aspergillus flaves*, *Tricothecium roseum* and *Aspergillus nigerr* shows high moderate disease severity(58.3%, 56.4% and 53.6%) respectively. This study curves creates another option for the management problem of *Eichhornia crassipes* in lake Tana by the use of pathogenic species as a biocontrol agents.

Key phrases: Biological control, pathogens, Disease incidence, Disease severity, Detached leaf test

